

ON INTEGRAL OPERATORS WITH OPERATOR VALUED KERNELS

RISHAD SHAHMUROV

ABSTRACT. Here $L_q \rightarrow L_p$ boundedness of integral operator with operator-valued kernels is studied and the main result is applied to convolution operators. Using these results $B_{q,r}^s \rightarrow B_{p,r}^s$ regularity for Fourier multiplier operator is established.

1. INTRODUCTION

It is well-known that, solutions of inhomogeneous differential and integral equations are represented by integral operators. In order to investigate stability properties of these problems it is important to have boundedness of corresponding integral operators in the studied function spaces. For instance, the boundedness of Fourier multiplier operators play crucial role in the theory of linear PDE especially in the study of maximal regularity for elliptic and parabolic PDE. For exposition of the integral operators with scalar valued kernels see [3] and for the application of multiplier theorems see [2].

Maria Girardi and Lutz Weis [4] recently proved that integral operator

$$(Kf)(\cdot) = \int_S k(\cdot, s)f(s)d\nu(s) \quad (1)$$

defines a bounded linear operator

$$K : L_p(S, X) \rightarrow L_p(T, Y)$$

provided some measurability conditions and the following assumptions

$$\begin{aligned} \sup_{s \in S} \int_T \|k(t, s)x\|_Y d\mu(t) &\leq C_1 \|x\|_X \text{ for all } x \in X, \\ \sup_{t \in T} \int_S \|k^*(t, s)y^*\|_{X^*} d\nu(s) &\leq C_2 \|y^*\|_{Y^*} \text{ for all } y^* \in Y^* \end{aligned}$$

are satisfied. Inspired from [4] we will show that (1) defines a bounded linear operator

$$K : L_q(S, X) \rightarrow L_p(T, Y)$$

if the kernel $k : T \times S \rightarrow B(X, Y)$ satisfies the following conditions

$$\sup_{s \in S} \left(\int_T \|k(t, s)x\|_Y^\theta dt \right)^{\frac{1}{\theta}} \leq C_1 \|x\|_X \text{ for all } x \in X,$$

2000 *Mathematics Subject Classification.* Primary 45P05, 47G10, 46E40 .

Key words and phrases. Integral operators, Banach-valued Bochner spaces, operator-valued kernels, interpolation of Banach spaces.

$$\sup_{t \in T} \left(\int_S \|k^*(t, s)y^*\|_{X^*}^\theta ds \right)^{\frac{1}{\theta}} \leq C_2 \|y^*\|_{Y^*} \text{ for all } y^* \in Y^*$$

where

$$\frac{1}{q} - \frac{1}{p} = 1 - \frac{1}{\theta}$$

for $1 \leq q < \frac{\theta}{\theta-1} \leq \infty$ and $\theta \in [1, \infty)$.

Here X and Y are Banach spaces over the field C and X^* is the dual space of X . The space $B(X, Y)$ of bounded linear operators from X to Y is endowed with the usual uniform operator topology.

Now let us state some important notations from [4]. A subspace Y of X^* τ -norms X , where $\tau \geq 1$, provided

$$\|x\|_X \leq \tau \sup_{x^* \in B(Y)} |x^*(x)| \quad \forall x \in X.$$

It is clear that if Y τ -norms X then the canonical mapping

$$u : X \rightarrow Y^* \text{ with } \langle y, ux \rangle = \langle x, y \rangle$$

is an isomorphic embedding with

$$\frac{1}{\tau} \|x\|_X \leq \|u(x)\|_{Y^*} \leq \|x\|_X.$$

Let $(T, \sum_T, \mu)$ and $(S, \sum_S, \nu)$ be positive measure spaces and

$$\sum_S^{\text{finite}} = \left\{ A \in \sum_S : \nu(A) < \infty \right\}, \quad \sum_S^{\text{full}} = \left\{ A \in \sum_S : \nu(S \setminus A) = 0 \right\}.$$

$\varepsilon(S, X)$ will denote the space of finitely-valued and finitely supported measurable functions from S into X i.e.

$$\varepsilon(S, X) = \left\{ \sum_{i=1}^n x_i 1_{A_i} : x_i \in X, A_i \in \sum_S^{\text{finite}}, n \in \mathbb{N} \right\}.$$

Note that $\varepsilon(S, X)$ is norm dense in $L_p(S, X)$ for $1 \leq p < \infty$. Let $L_\infty^0(S, X)$ be the closure of $\varepsilon(S, X)$ in $L_\infty(S, X)$ norm. In general $L_\infty^0(S, X) \neq L_\infty(S, X)$ (see [4, Proposition 2.2] and [4, Lemma 2.3]).

A vector valued function $f : S \rightarrow X$ is measurable if there is a sequence $(f_n)_{n=1}^\infty \subset \varepsilon(S, X)$ converging (in the sense of X topology) to f and it is $\sigma(X, \Gamma)$ -measurable provided $\langle f(\cdot), x^* \rangle : S \rightarrow K$ is measurable for each $x^* \in \Gamma \subset X^*$. Suppose $1 \leq p \leq \infty$ and $\frac{1}{p} + \frac{1}{p'} = 1$. There is a natural isometric embedding of $L_{p'}(T, Y^*)$ into $[L_p(T, Y)]^*$ given by

$$\langle f, g \rangle = \int_T \langle f(t), g(t) \rangle d\mu(t) \text{ for } g \in L_{p'}(T, Y^*) \text{ and } f \in L_p(T, Y).$$

Often $[E(X)]^* = E^*(X^*)$, for example, provided X has the Radon-Nikodym property. Let us remind an important fact that if X is reflexive or if X is separable, then X has the Radon-Nikodym property.

2. $L_q \rightarrow L_p$ ESTIMATES FOR INTEGRAL OPERATORS

In this section we identify conditions on operator-valued kernel $k : T \times S \rightarrow B(X, Y)$, extending theorems in [4] so that

$$\|K\|_{L_q(S, X) \rightarrow L_p(T, Y)} \leq C$$

for $1 \leq q \leq p$. To prove our main result we shall use interpolation theorems of L_p spaces. Therefore, we will study $L_1(S, X) \rightarrow L_\theta(T, Y)$ and $L_{\theta'}(S, X) \rightarrow L_\infty(T, Y)$ boundedness of integral operator (1). The following two conditions are natural measurability assumptions on $k : T \times S \rightarrow B(X, Y)$.

Condition 2.1. For any $A \in \sum_S^{\text{finite}}$ and each $x \in X$

(a) there is $T_{A,x} \in \sum_T^{\text{full}}$ so that if $t \in T_{A,x}$ then the Bochner integral

$$\int_A k(t, s)x d\nu(s) \text{ exists,}$$

(b) $T_{A,x} : t \rightarrow \int_A k(t, s)x d\nu(s)$ defines a measurable function from T into Y .

Note that if k satisfies the above condition then for each $f \in \varepsilon(S, X)$, there is $T_f \in \sum_T^{\text{full}}$ so that the Bochner integral

$$\int_S k(t, s)f(s)d\nu(s) \text{ exists}$$

and (1) defines a linear mapping

$$K : \varepsilon(S, X) \rightarrow L_0(T, Y)$$

where L_0 denotes the space of measurable functions.

Condition 2.2. The kernel $k : T \times S \rightarrow B(X, Y)$ satisfies following properties:

(a) a real valued mapping $\|k(t, s)x\|_X^\theta$ is product measurable $\forall x \in X$

(b) there is $S_x \in \sum_S^{\text{full}}$ so that

$$\|k(t, s)x\|_{L_\theta(T, Y)} \leq C_1 \|x\|_X$$

for $1 \leq \theta < \infty$ and $x \in X$.

Theorem 2.3. Suppose $1 \leq \theta < \infty$ and the kernel $k : T \times S \rightarrow B(X, Y)$ satisfies Condition 2.1 and Condition 2.2. Then integral operator (1) acting on $\varepsilon(S, X)$ extends to a bounded linear operator

$$K : L_1(S, X) \rightarrow L_\theta(T, Y).$$

Proof. Let $f = \sum_{i=1}^n x_i 1_{A_i}(s) \in \varepsilon(S, X)$ be fixed. Taking into account the fact that $1 \leq \theta$, using the general Minkowski-Jessen inequality and assumptions of the

theorem we obtain

$$\begin{aligned}
\|(Kf)(t)\|_{L_\theta(T,Y)} &\leq \left[\int_T \left(\int_S \|k(t,s) \sum_{i=1}^n x_i 1_{A_i}(s)\|_Y d\nu(s) \right)^\theta d\mu(t) \right]^{\frac{1}{\theta}} \\
&\leq \int_S \left(\int_T \|k(t,s) \sum_{i=1}^n x_i 1_{A_i}(s)\|_Y^\theta d\mu(t) \right)^{\frac{1}{\theta}} d\nu(s) \\
&\leq \int_S \left[\int_T \left(\sum_{i=1}^n 1_{A_i}(s) \|k(t,s)x_i\|_Y \right)^\theta d\mu(t) \right]^{\frac{1}{\theta}} d\nu(s) \\
&\leq \int_S \sum_{i=1}^n 1_{A_i}(s) \left(\int_T \|k(t,s)x_i\|_Y^\theta d\mu(t) \right)^{\frac{1}{\theta}} d\nu(s) \\
&\leq \int_S \sum_{i=1}^n 1_{A_i}(s) \|k(t,s)x_i\|_{L_\theta(T,Y)} d\nu(s) \leq C_1 \sum_{i=1}^n \|x_i\|_X \int_S 1_{A_i}(s) d\nu(s) \\
&= C_1 \sum_{i=1}^n \|x_i\|_X \nu(A_i) = C_1 \|f\|_{L_1(S,X)}.
\end{aligned}$$

Hence, $\|K\|_{L_1 \rightarrow L_\theta} \leq C_1$. ■

Condition 2.4. For each $y^* \in Z$ there is $T_{y^*} \in \sum_T^{\text{full}}$ so that $\forall t \in T_{y^*}$

(a) a real valued mapping $\|k^*(t,s)x^*\|_{X^*}^\theta$ is measurable $\forall x^* \in X^*$

(b) There is $S_x \in \sum_S^{\text{full}}$ so that

$$\|k^*(t,s)y^*\|_{L_\theta(S,X^*)} \leq C_2 \|y^*\|_{Y^*}$$

for $1 \leq \theta < \infty$ and $x \in X$.

Theorem 2.5. Let Z be a subspace of Y^* that τ -norms Y . Suppose $1 \leq \theta < \infty$ and $k : T \times S \rightarrow B(X,Y)$ satisfies Condition 2.1 and Condition 2.4. Then integral operator (1) acting on $\varepsilon(S,X)$ extends to a bounded linear operator

$$K : L_{\theta'}(S,X) \rightarrow L_\infty(T,Y).$$

Proof. Suppose $f \in \varepsilon(S,X)$ and $y^* \in Z$ are fixed. Let $T_f, T_{y^*} \in \sum_T^{\text{full}}$ be corresponding sets due to Condition 2.1 and Condition 2.4. If $t \in T_f \cap T_{y^*}$ then by using Hölder's inequality and assumptions of the theorem we get

$$\begin{aligned}
| \langle y^*, (Kf)(t) \rangle_Y | &= \left| \left\langle y^*, \int_S k(t,s)f(s) d\nu(s) \right\rangle \right| \\
&\leq \int_S |[k^*(t,s)y^*] f(s)| d\nu(s) \\
&\leq \|k^*(t,s)y^*\|_{L_\theta(S,X^*)} \|f(s)\|_{L_{\theta'}(S,X)} \\
&\leq C_2 \|y^*\| \|f\|_{L_{\theta'}(S,X)}.
\end{aligned}$$

Since, $T_f \cap T_{y^*} \in \sum_T^{\text{full}}$ and Z τ -norms Y

$$\|Kf\|_{L_\infty(T,Y)} \leq C_2 \tau \|f\|_{L_{\theta'}(S,X)}.$$

Hence, $\|K\|_{L_{\theta'} \rightarrow L_\infty} \leq \tau C_2$. ■

In [4, Lemma 3.9] authors slightly improved interpolation theorem [1, Thm 5.1.2]. The next lemma is a more general form of [4, Lemma 3.9].

Lemma 2.6. Suppose a linear operator

$$K : \varepsilon(S, X) \rightarrow L_\theta(T, Y) + L_\infty(T, Y)$$

satisfies

$$\|Kf\|_{L_\theta(T,Y)} \leq C_1 \|f\|_{L_1(S,X)} \text{ and } \|Kf\|_{L_\infty(T,Y)} \leq C_2 \|f\|_{L_{\theta'}(S,X)} \quad (2)$$

Then, for $\frac{1}{q} - \frac{1}{p} = 1 - \frac{1}{\theta}$ and $1 \leq q < \frac{\theta}{\theta-1} \leq \infty$ the mapping K extends to a bounded linear operator

$$K : L_q(S, X) \rightarrow L_p(T, Y)$$

with

$$\|K\|_{L_q \rightarrow L_p} \leq (C_1)^{\frac{\theta}{p}} (C_2)^{1-\frac{\theta}{p}}.$$

Proof. Let us first consider conditional expectation operator

$$(K_0 f) = E((Kf)_{1_B} | \Sigma)$$

where Σ is a σ -algebra of subsets of $B \in \sum_T^{\text{finite}}$. From (2) it follows that

$$\begin{aligned} \|K_0 f\|_{L_\theta(T,Y)} &\leq C_1 \|f\|_{L_1(S,X)} < \infty \text{ and} \\ \|K_0 f\|_{L_\infty(T,Y)} &\leq C_2 \|f\|_{L_{\theta'}(S,X)} < \infty. \end{aligned}$$

Hence, by Riesz-Thorin theorem ([1, Thm 5.1.2]) we have

$$\|K_0 f\|_{L_p(T,Y)} \leq (C_1)^{\frac{\theta}{p}} (C_2)^{1-\frac{\theta}{p}} \|f\|_{L_q(S,X)}. \quad (3)$$

Now, taking into account (3) and using the same reasoning as in the proof of [4, Lemma 3.9] one can easily show assertion of this lemma. ■

Theorem 2.7. (Operator-valued Schur's test) Let Z be a subspace of Y^* that τ -norms Y and $\frac{1}{q} - \frac{1}{p} = 1 - \frac{1}{\theta}$ for $1 \leq q < \frac{\theta}{\theta-1} \leq \infty$. Suppose $k : T \times S \rightarrow B(X, Y)$ satisfies Condition 2.1, Condition 2.2 and Condition 2.4 with respect to Z . Then integral operator (1) extends to a bounded linear operator

$$K : L_q(S, X) \rightarrow L_p(T, Y)$$

with

$$\|K\|_{L_q \rightarrow L_p} \leq (C_1)^{\frac{\theta}{p}} (\tau C_2)^{1-\frac{\theta}{p}}.$$

Proof. Combining Theorem 2.3, Theorem 2.5 and Lemma 2.6 we obtain assertion of the theorem. ■

Remark 2.8. Note that choosing $\theta = 1$ we get original results in [4].

For L_∞ estimates (it is more delicate and based on ideas from geometry Banach spaces) and weak continuity and duality results see [4]. The next corollary plays important role in the Fourier Multiplier theorems.

Corollary 2.9. Let Z be a subspace of Y^* that τ -norms Y and $\frac{1}{q} - \frac{1}{p} = 1 - \frac{1}{\theta}$ for $1 \leq q < \frac{\theta}{\theta-1} \leq \infty$. Suppose $k : R^n \rightarrow B(X, Y)$ is strongly measurable on X , $k^* : R^n \rightarrow B(Y^*, X^*)$ is strongly measurable on Z and

$$\begin{aligned} \|kx\|_{L_\theta(R^n, Y)} &\leq C_1 \|x\|_X \text{ for all } x \in X, \\ \|k^*y^*\|_{L_\theta(R^n, X^*)} &\leq C_2 \|y^*\|_{Y^*} \text{ for all } y^* \in Y^* \end{aligned}$$

Then the convolution operator defined by

$$(Kf)(t) = \int_{R^n} k(t-s)f(s)ds \text{ for } t \in R^n$$

satisfies $\|K\|_{L_q \rightarrow L_p} \leq (C_1)^{\frac{\theta}{p}} (C_2)^{1-\frac{\theta}{p}}$.

It is easy to see that $k : R^n \rightarrow B(X, Y)$ satisfies Condition 2.1, Condition 2.2 and Condition 2.4 with respect to Z . Thus, assertion of corollary follows from Theorem 2.7.

3. FOURIER MULTIPLIERS OF BESOV SPACES

In this section we shall indicate importance of Corollary 2.9 in the theory of Fourier multipliers (FM). Thus we give definition and some basic properties of operator valued FM and Besov spaces.

Consider some subsets $\{J_k\}_{k=0}^\infty$ and $\{I_k\}_{k=0}^\infty$ of R^n , where

$$J_0 = \{t \in R^n : |t| \leq 1\}, \quad J_k = \{t \in R^n : 2^{k-1} \leq |t| \leq 2^k\} \text{ for } k \in N$$

and

$$I_0 = \{t \in R^n : |t| \leq 2\}, \quad I_k = \{t \in R^n : 2^{k-1} \leq |t| \leq 2^{k+1}\} \text{ for } k \in N.$$

Let us define the partition of unity $\{\varphi_k\}_{k \in N_0}$ of functions from $S(R^n, R)$. Suppose $\psi \in S(R, R)$ is a nonnegative function with support in $[2^{-1}, 2]$, which satisfies

$$\sum_{k=-\infty}^{\infty} \psi(2^{-k}s) = 1 \text{ for } s \in R \setminus \{0\}$$

and

$$\varphi_k(t) = \psi(2^{-k}|t|), \quad \varphi_0(t) = 1 - \sum_{k=1}^{\infty} \varphi_k(t) \text{ for } t \in R^n.$$

Let $1 \leq q \leq r \leq \infty$ and $s \in R$. The Besov space is the set of all functions $f \in S'(R^n, X)$ for which

$$\begin{aligned} \|f\|_{B_{q,r}^s(R^n, X)} : &= \|2^{ks} \{(\tilde{\varphi}_k * f)\}_{k=0}^\infty\|_{l_r(L_q(R^n, X))} \\ &\equiv \begin{cases} \left[\sum_{k=0}^{\infty} 2^{ksr} \|\tilde{\varphi}_k * f\|_{L_q(R^n, X)}^r \right]^{\frac{1}{r}} & \text{if } r \neq \infty \\ \sup_{k \in N_0} [2^{ks} \|\tilde{\varphi}_k * f\|_{L_q(R^n, X)}] & \text{if } r = \infty \end{cases} \end{aligned}$$

is finite; here q and s are main and smoothness indexes respectively. The Besov space has significant interpolation and embedding properties:

$$B_{q,r}^s(R^n; X) = (L_q(R^n; X), W_q^m(R^d; X))_{\frac{s}{m}, r},$$

$$W_q^{l+1}(X) \hookrightarrow B_{q,r}^s(X) \hookrightarrow W_q^l(X) \hookrightarrow L_q(X) \text{ where } l < s < l+1,$$

$$B_{\infty,1}^s(X) \hookrightarrow C^s(X) \hookrightarrow B_{\infty,\infty}^s(X) \text{ for } s \in \mathbf{Z},$$

and

$$B_{p,1}^{\frac{d}{p}}(R^d, X) \hookrightarrow L_\infty(R^d, X) \text{ for } s \in \mathbf{Z},$$

where $m \in N$ and $C^s(X)$ denotes the Holder-Zygmund spaces.

Definition 3.1. Let X be a Banach space and $1 \leq u \leq 2$. We say X has Fourier type u if

$$\|\mathcal{F}f\|_{L_{u'}(R^n, X)} \leq C\|f\|_{L_u(R^n, X)} \text{ for each } f \in S(R^n, X),$$

where $\frac{1}{u} + \frac{1}{u'} = 1$, $\mathcal{F}_{u,n}(X)$ is the smallest $C \in [0, \infty]$. Let us list some important facts:

- (i) Any Banach space has a Fourier type 1,
- (ii) B -convex Banach spaces have a nontrivial Fourier type,
- (iii) Spaces having Fourier type 2 should be isomorphic to a Hilbert spaces.

From [5, **Theorem 3.1**] it follows the following corollary:

Corollary 3.2. Let X be a Banach space having Fourier type $u \in [1, 2]$ and $1 \leq \theta \leq u'$. Then the inverse Fourier transform defines a bounded operator

$$\mathcal{F}^{-1} : B_{u,1}^{n(\frac{1}{\theta} - \frac{1}{u'})}(R^n, X) \rightarrow L_\theta(R^n, X). \quad (4)$$

Definition 3.3. Let $(E_1(R^n, X), E_2(R^n, Y))$ be one of the following systems, where $1 \leq q \leq p \leq \infty$

$$(L_q(X), L_p(Y)) \text{ or } (B_{q,r}^s(X), B_{p,r}^s(Y)).$$

A bounded measurable function $m : R^n \rightarrow B(X, Y)$ is called a Fourier multiplier from $E_1(X)$ to $E_2(Y)$ if there is a bounded linear operator

$$T_m : E_1(X) \rightarrow E_2(Y)$$

such that

$$T_m(f) = \mathcal{F}^{-1}[m(\cdot)(\mathcal{F}f)(\cdot)] \text{ for each } f \in S(X), \quad (5)$$

$$T_m \text{ is } \sigma(E_1(X), E_1^*(X^*)) \text{ to } \sigma(E_2(Y), E_2^*(Y^*)) \text{ continuous.} \quad (6)$$

The uniquely determined operator T_m is the FM operator induced by m . Note that if $T_m \in B(E_1(X), E_2(Y))$ and T_m^* maps $E_2^*(Y^*)$ into $E_1^*(X^*)$ then T_m satisfies the weak continuity condition (6).

For definition of Besov spaces and its basic properties we refer to [5].

Since (5) can be written in the convolution form

$$T_m(f)(t) = \int_{R^n} \check{m}(t-s)f(s)ds \quad (7)$$

Corollary 2.9 and Corollary 3.2 can be applied to obtain $L_q(R^n, X) \rightarrow L_p(R^n, Y)$ regularity for (5).

Theorem 3.4. Let X and Y be Banach spaces having Fourier type $u \in [1, 2]$ and $p, q \in [1, \infty]$ so that $0 \leq \frac{1}{q} - \frac{1}{p} \leq \frac{1}{u}$. Then there is a constant C depending only on $\mathcal{F}_{u,n}(X)$ and $\mathcal{F}_{u,n}(Y)$ so that if

$$m \in B_{u,1}^{n(\frac{1}{u} + \frac{1}{p} - \frac{1}{q})}(R^n, B(X, Y))$$

then m is a FM from $L_q(R^n, X)$ to $L_p(R^n, Y)$ with

$$\|T_m\|_{L_q(R^n, X) \rightarrow L_p(R^n, Y)} \leq CM_u(m) \text{ for each}$$

where

$$M_u(m) = \inf \left\{ \|m(a \cdot)\|_{B_{u,1}^{n(\frac{1}{u} + \frac{1}{p} - \frac{1}{q})}(R^n, B(X, Y))} : a > 0 \right\}.$$

Proof. Let $\frac{1}{q} - \frac{1}{p} = 1 - \frac{1}{\theta}$ and $1 \leq q < \frac{\theta}{\theta-1} \leq \infty$. Assume that $m \in S(B(X, Y))$. Then $\tilde{m} \in S(B(X, Y))$. Since $\mathcal{F}^{-1}[m(a \cdot)x](s) = a^{-n} \tilde{m}(\frac{s}{a})x$, choosing appropriate a and using (4) we obtain

$$\begin{aligned} \|\tilde{m}x\|_{L_\theta(Y)} &= \|[m(a \cdot)x]^\vee\|_{L_\theta(Y)} \\ &\leq C_1 \|m(a \cdot)\|_{B_{u,1}^{n(\frac{1}{\theta} - \frac{1}{u})}} \|x\|_X \\ &\leq 2C_1 M_u(m) \|x\|_X \end{aligned} \quad (8)$$

where C_1 depends only on $\mathcal{F}_{u,n}(Y)$. Since $m \in S(B(X, Y))$ we have $[m^*]^\vee = [\tilde{m}]^* \in S(B(Y^*, X^*))$ and $M_u(m) = M_u(m^*)$. Thus, in a similar manner as above, we get

$$\|[\tilde{m}(\cdot)]^* y^*\|_{L_\theta(Y)} \leq 2C_2 M_u(m) \|y^*\|_{Y^*} \quad (9)$$

for some constant C_2 depending on $\mathcal{F}_{u,n}(X^*)$. Hence by (8-9) and Corollary 2.9

$$(T_m f)(t) = \int_{R^n} \tilde{m}(t-s) f(s) ds$$

satisfies

$$\|T_m f\|_{L_p(R^n, Y)} \leq CM_u(m) \|f\|_{L_q(R^n, X)}$$

for all $p, q \in [1, \infty]$ so that $0 \leq \frac{1}{q} - \frac{1}{p} \leq \frac{1}{u}$. Now, taking into account the fact that $S(B(X, Y))$ is continuously embedded in $B_{u,1}^{n(\frac{1}{u} + \frac{1}{p} - \frac{1}{q})}(B(X, Y))$ and using the same reasoning as [5, Theorem 4.3] one can easily prove the general case $m \in B_{u,1}^{n(\frac{1}{u} + \frac{1}{p} - \frac{1}{q})}$ and weak continuity of T_m . ■

Theorem 3.5. Let X and Y be Banach spaces having Fourier type $u \in [1, 2]$ and $p, q \in [1, \infty]$ are so that $0 \leq \frac{1}{q} - \frac{1}{p} \leq \frac{1}{u}$. Then, there exist a constant C depending only on $\mathcal{F}_{u,n}(X)$ and $\mathcal{F}_{u,n}(Y)$ so that if $m : R^n \rightarrow B(X, Y)$ satisfy

$$\varphi_k \cdot m \in B_{u,1}^{n(\frac{1}{u} + \frac{1}{p} - \frac{1}{q})}(R^n, B(X, Y)) \text{ and } M_u(\varphi_k \cdot m) \leq A \quad (10)$$

then m is a FM from $B_{q,r}^s(R^n, X)$ to $B_{p,r}^s(R^n, Y)$ and $\|T_m\|_{B_{q,r}^s \rightarrow B_{p,r}^s} \leq CA$ for each $s \in R$ and $r \in [1, \infty]$.

Taking into consideration the Theorem 3.4 one can easily prove above theorem in a similar manner as [5, Theorem 4.3].

The following corollary provides a practical sufficient condition to check (10).

Lemma 3.6. Let $n \left(\frac{1}{u} + \frac{1}{p} - \frac{1}{q} \right) < l \in N$ and $\theta \in [u, \infty]$. If $m \in C^l(R^n, B(X, Y))$ and

$$\|D^\alpha m|_{I_0}\|_{L_\theta(B(X, Y))} \leq A, \quad \|D^\alpha m_k|_{I_1}\|_{L_\theta(B(X, Y))} \leq A, \quad m_k(\cdot) = m(2^{k-1} \cdot),$$

for each $\alpha \in N^n$, $|\alpha| \leq l$ and $k \in N$ then m satisfies (10).

Using the fact that $W_u^l(R^n, B(X, Y)) \subset B_{u,1}^{n(\frac{1}{u} + \frac{1}{p} - \frac{1}{q})}(R^n, B(X, Y))$, the above lemma can be proven in similar fashion as [5, Lemma 4.10].

Choosing $\theta = \infty$ in the Lemma 3.6 we get the following corollary:

Corollary 3.7. (Mikhlin's condition) Let X and Y be Banach spaces having Fourier type $u \in [1, 2]$ and $0 \leq \frac{1}{q} - \frac{1}{p} \leq \frac{1}{u}$. If $m \in C^l(R^n, B(X, Y))$ satisfies

$$\left\| (1 + |t|)^{|\alpha|} D^\alpha m(t) \right\|_{L^\infty(R^n, B(X, Y))} \leq A$$

for each multi-index α with $|\alpha| \leq l = \left\lceil n \left(\frac{1}{u} + \frac{1}{p} - \frac{1}{q} \right) \right\rceil + 1$, then m is a FM from $B_{q,r}^s(R^n, X)$ to $B_{p,r}^s(R^n, Y)$ for each $s \in \mathbb{R}$ and $r \in [1, \infty]$.

Remark 3.8. Corollary 2.13 particularly implies the following facts:

- (a) If X and Y are arbitrary Banach spaces then $l = \left\lceil n \left(\frac{1}{p} + \frac{1}{q'} \right) \right\rceil + 1$,
- (b) If X and Y be Banach spaces having Fourier type $u \in [1, 2]$ and $\frac{1}{q} - \frac{1}{p} = \frac{1}{u}$ then Miklin's condition with order 1 suffices for multiplier function to be a FM in $(B_{q,r}^s(R^n, X), B_{p,r}^s(R^n, Y))$.
- (c) Let X and Y be Hilbert spaces and $p, q \in [1, \infty]$ are so that $\frac{1}{q} - \frac{1}{p} = \frac{1}{2}$. If $m \in C^1(R^n, B(X, Y))$ satisfies

$$\sup_{t \in R^n} \|m(t)\|_{B(X, Y)} \leq A_1,$$

$$\sup_{t \in R^n} \left\| (1 + |t|) \frac{d}{dt} m(t) \right\|_{B(X, Y)} \leq A_2$$

then m is a FM from $B_{q,r}^s(R^n, X)$ to $B_{p,r}^s(R^n, Y)$.

REFERENCES

- [1] J. B rgh, J. L fstr m, *Interpolation spaces: An introduction*, Springer-Verlag, Berlin 1976, Grundlehren der Mathematischen Wissenschaften, No. 223.
- [2] R. Denk, M. Hieber, J. Pr uss, R -boundedness, Fourier multipliers and problems of elliptic and parabolic type, *Mem. Amer. Math. Soc.* 166(2003), p. 1-106.
- [3] G. Folland, *Real analysis, Pure and Applied Mathematics*, John Wiley & Sons Inc., New York, 1984, Modern techniques and their applications, A Wiley-Interscience Publication.
- [4] M. Girardi, L. Weis, Integral operators with operator-valued kernels, *Journal of Mathematical Analysis and Applications*, 290 (2004), no.1, pp.190-212.
- [5] M. Girardi, L. Weis, Operator-valued multiplier theorems on Besov spaces, *Math. Nachr.*, **251** (2003), 34-51.

DEPARTMENT OF MATHEMATICS, YEDITEPE UNIVERSITY, KAYISHDAGI CADDESİ, 34755 KAYISHDAGI, ISTANBUL, TURKEY

E-mail address: shahmurov@hotmail.com